

Moment Varieties of Measures on Polytopes

joint works with Kristian Ranestad (Universitetet i Oslo),
Boris Shapiro (Stockholms universitet)
and Bernd Sturmfels (UC Berkeley / MPI MiS Leipzig)

July 10, 2019

Moments of a Polytope

- ◆ Let $P \subset \mathbb{R}^d$ be a full-dimensional polytope.
- ◆ μ_P : uniform probability distribution on P
- ◆ moments

$$m_{i_1 i_2 \dots i_d}(P) := \int_{\mathbb{R}^d} w_1^{i_1} w_2^{i_2} \dots w_d^{i_d} d\mu_P \quad \text{for } i_1, i_2, \dots, i_d \in \mathbb{Z}_{\geq 0}$$

Moments of a Polytope

- ◆ Let $P \subset \mathbb{R}^d$ be a full-dimensional polytope.
- ◆ μ_P : uniform probability distribution on P
- ◆ moments

$$m_{i_1 i_2 \dots i_d}(P) := \int_{\mathbb{R}^d} w_1^{i_1} w_2^{i_2} \dots w_d^{i_d} d\mu_P \quad \text{for } i_1, i_2, \dots, i_d \in \mathbb{Z}_{\geq 0}$$

Known:

The list of all moments $(m_{\mathcal{I}}(P) \mid \mathcal{I} \in \mathbb{Z}_{\geq 0}^d)$ uniquely encodes P .

↪ Can recover P from its moments.

Moments of a Polytope

- ◆ Let $P \subset \mathbb{R}^d$ be a full-dimensional polytope.
- ◆ μ_P : uniform probability distribution on P
- ◆ moments

$$m_{i_1 i_2 \dots i_d}(P) := \int_{\mathbb{R}^d} w_1^{i_1} w_2^{i_2} \dots w_d^{i_d} d\mu_P \quad \text{for } i_1, i_2, \dots, i_d \in \mathbb{Z}_{\geq 0}$$

Known:

The list of all moments $(m_{\mathcal{I}}(P) \mid \mathcal{I} \in \mathbb{Z}_{\geq 0}^d)$ uniquely encodes P .

↪ Can recover P from its moments.

Caution: The moments are not independent of each other.

Our Goal:

Study the dependencies among the moments!

Moment Varieties

- ◆ We assume: $P \subset \mathbb{R}^d$ is full-dimensional, simplicial, and has n vertices

Moment Varieties

- ◆ We assume: $P \subset \mathbb{R}^d$ is full-dimensional, simplicial, and has n vertices
- ◆ We can vary the vertices of P locally without changing the combinatorial type \mathcal{P} of P
 - $\rightsquigarrow m_{\mathcal{I}}(P)$ is a locally defined function of the vertex coordinates

Moment Varieties

- ◆ We assume: $P \subset \mathbb{R}^d$ is full-dimensional, simplicial, and has n vertices
- ◆ We can vary the vertices of P locally without changing the combinatorial type \mathcal{P} of P
- ↪ $m_{\mathcal{I}}(P)$ is a locally defined function of the vertex coordinates
- ↪ For every combinatorial type \mathcal{P} and every finite subset $\mathcal{A} \subset \mathbb{Z}_{\geq 0}^d$, we have a rational function

$$m_{\mathcal{P}, \mathcal{A}} : (\mathbb{R}^d)^n \dashrightarrow \mathbb{R}^{|\mathcal{A}|},$$
$$P \longmapsto (m_{\mathcal{I}}(P))_{\mathcal{I} \in \mathcal{A}}$$

Moment Varieties

- ◆ We assume: $P \subset \mathbb{R}^d$ is full-dimensional, simplicial, and has n vertices
- ◆ We can vary the vertices of P locally without changing the combinatorial type \mathcal{P} of P
 - $\rightsquigarrow m_{\mathcal{I}}(P)$ is a locally defined function of the vertex coordinates
 - \rightsquigarrow For every combinatorial type \mathcal{P} and every finite subset $\mathcal{A} \subset \mathbb{Z}_{\geq 0}^d$, we have a rational function

$$m_{\mathcal{P}, \mathcal{A}} : (\mathbb{R}^d)^n \dashrightarrow \mathbb{R}^{|\mathcal{A}|},$$
$$P \longmapsto (m_{\mathcal{I}}(P))_{\mathcal{I} \in \mathcal{A}}$$

- ◆ We assume: $0 \in \mathcal{A} \rightsquigarrow m_{\mathcal{P}, \mathcal{A}} : \mathbb{C}^{d \times n} \dashrightarrow \mathbb{P}_{\mathbb{C}}^{|\mathcal{A}|-1}$

Moment Varieties

- ◆ We assume: $P \subset \mathbb{R}^d$ is full-dimensional, simplicial, and has n vertices
- ◆ We can vary the vertices of P locally without changing the combinatorial type \mathcal{P} of P
 - $\rightsquigarrow m_{\mathcal{I}}(P)$ is a locally defined function of the vertex coordinates
 - \rightsquigarrow For every combinatorial type \mathcal{P} and every finite subset $\mathcal{A} \subset \mathbb{Z}_{\geq 0}^d$, we have a rational function

$$m_{\mathcal{P}, \mathcal{A}} : (\mathbb{R}^d)^n \dashrightarrow \mathbb{R}^{|\mathcal{A}|},$$
$$P \longmapsto (m_{\mathcal{I}}(P))_{\mathcal{I} \in \mathcal{A}}$$

- ◆ We assume: $0 \in \mathcal{A} \rightsquigarrow m_{\mathcal{P}, \mathcal{A}} : \mathbb{C}^{d \times n} \dashrightarrow \mathbb{P}_{\mathbb{C}}^{|\mathcal{A}|-1}$
- ◆ **Moment variety**

$$\mathcal{M}_{\mathcal{A}}(\mathcal{P}) := \overline{m_{\mathcal{P}, \mathcal{A}}(\mathbb{C}^{d \times n})} \subset \mathbb{P}_{\mathbb{C}}^{|\mathcal{A}|-1}$$

Example: Line Segments

◆ Let $P = [a, b] \subset \mathbb{R}^1$

$$\begin{aligned}\Rightarrow m_i(P) = m_i(a, b) &= \frac{1}{b-a} \int_a^b w^i dw = \frac{1}{i+1} \frac{b^{i+1} - a^{i+1}}{b-a} \\ &= \frac{1}{i+1} (a^i + a^{i-1}b + a^{i-2}b^2 + \dots + b^i)\end{aligned}$$

Example: Line Segments

◆ Let $P = [a, b] \subset \mathbb{R}^1$

$$\begin{aligned}\Rightarrow m_i(P) &= m_i(a, b) = \frac{1}{b-a} \int_a^b w^i dw = \frac{1}{i+1} \frac{b^{i+1} - a^{i+1}}{b-a} \\ &= \frac{1}{i+1} (a^i + a^{i-1}b + a^{i-2}b^2 + \dots + b^i)\end{aligned}$$

$\Rightarrow m_{\text{LineSegments}, \{0, 1, \dots, r\}} : \mathbb{C}^2 \dashrightarrow \mathbb{P}^r,$

$$(a, b) \longmapsto (m_0(a, b) : m_1(a, b) : \dots : m_r(a, b))$$

Example: Line Segments


- ◆ Let $P = [a, b] \subset \mathbb{R}^1$

$$\begin{aligned}\Rightarrow m_i(P) &= m_i(a, b) = \frac{1}{b-a} \int_a^b w^i dw = \frac{1}{i+1} \frac{b^{i+1} - a^{i+1}}{b-a} \\ &= \frac{1}{i+1} (a^i + a^{i-1}b + a^{i-2}b^2 + \dots + b^i)\end{aligned}$$

$$\begin{aligned}\Rightarrow m_{\text{LineSegments}, \{0,1,\dots,r\}} : \mathbb{C}^2 &\dashrightarrow \mathbb{P}^r, \\ (a, b) &\longmapsto (m_0(a, b) : m_1(a, b) : \dots : m_r(a, b))\end{aligned}$$

- ◆ $\mathcal{M}_{\{0,1,\dots,r\}}(\text{LineSegments})$ is a surface in \mathbb{P}^r


Example: Line Segments



Moment surface $\mathcal{M}_{\{0,1,2,3\}}(\text{LineSegments}) \subset \mathbb{P}^3$ in affine chart $\{m_0 = 1\}$

- ◆ Defined by $2m_1^3 - 3m_0m_1m_2 + m_0^2m_3 = 0$


Example: Line Segments



Moment surface $\mathcal{M}_{\{0,1,2,3\}}(\text{LineSegments}) \subset \mathbb{P}^3$ in affine chart $\{m_0 = 1\}$

- ◆ Defined by $2m_1^3 - 3m_0m_1m_2 + m_0^2m_3 = 0$
- ◆ Singular along $\{m_0 = m_1 = 0\}$

Example: Line Segments




Moment surface $\mathcal{M}_{\{0,1,2,3\}}(\text{LineSegments}) \subset \mathbb{P}^3$ in affine chart $\{m_0 = 1\}$

- ◆ Defined by $2m_1^3 - 3m_0m_1m_2 + m_0^2m_3 = 0$
- ◆ Singular along $\{m_0 = m_1 = 0\}$
- ◆ Contains twisted cubic curve (in red) corresponding to degenerate line segments $[a, a]$ of length 0

Example: Quadrilaterals

Let $\mathcal{A} := \{\mathcal{I} \in \mathbb{Z}_{\geq 0}^2 \mid |\mathcal{I}| \leq 3\}$.


Can we compute the moment hypersurface $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$?



Example: Quadrilaterals

Let $\mathcal{A} := \{\mathcal{I} \in \mathbb{Z}_{\geq 0}^2 \mid |\mathcal{I}| \leq 3\}$.

Can we compute the moment hypersurface $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$?



Observations:

The defining equation of $\mathcal{M}_{\mathcal{A}}(\square)$ is


- ♦ homogeneous with respect to the \mathbb{Z}^3 -grading given by

$$\text{degree}(m_{i_1 i_2}) = (1, i_1, i_2),$$

Example: Quadrilaterals

Let $\mathcal{A} := \{\mathcal{I} \in \mathbb{Z}_{\geq 0}^2 \mid |\mathcal{I}| \leq 3\}$.

Can we compute the moment hypersurface $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$?



Observations:

The defining equation of $\mathcal{M}_{\mathcal{A}}(\square)$ is

- ◆ homogeneous with respect to the \mathbb{Z}^3 -grading given by

$$\text{degree}(m_{i_1 i_2}) = (1, i_1, i_2),$$

- ◆ invariant under the natural action of the affine group Aff_2 .

Example: Quadrilaterals

Using monodromy methods from numerical algebraic geometry, we compute that the defining equation of $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$ has \mathbb{Z}^3 -degree **(18, 12, 12)**.

Example: Quadrilaterals

Using monodromy methods from numerical algebraic geometry, we compute that the defining equation of $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$ has \mathbb{Z}^3 -degree **(18, 12, 12)**.

Proposition (K., Shapiro, Sturmfels)

The invariant ring $\mathbb{R}[m_{\mathcal{I}} \mid \mathcal{I} \in \mathcal{A}]^{\text{Aff}^2}$ is generated by **6 invariants**:

	m_{00}	s	t	h	g	j
\mathbb{Z}^3 -degree	(1, 0, 0)	(4, 4, 4)	(6, 6, 6)	(3, 2, 2)	(8, 6, 6)	(12, 9, 9)
# terms	1	25	103	5	168	892

Example: Quadrilaterals

Using monodromy methods from numerical algebraic geometry, we compute that the defining equation of $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$ has \mathbb{Z}^3 -degree **(18, 12, 12)**.

Proposition (K., Shapiro, Sturmfels)

The invariant ring $\mathbb{R}[m_{\mathcal{I}} \mid \mathcal{I} \in \mathcal{A}]^{\text{Aff}^2}$ is generated by **6 invariants**:

	m_{00}	s	t	h	g	j
\mathbb{Z}^3 -degree	(1, 0, 0)	(4, 4, 4)	(6, 6, 6)	(3, 2, 2)	(8, 6, 6)	(12, 9, 9)
# terms	1	25	103	5	168	892

We use the moments of various random quadrilaterals to **interpolate** the defining equation of $\mathcal{M}_{\mathcal{A}}(\square) \subset \mathbb{P}^9$:

$$\begin{aligned} & 2125764 h^6 + 5484996 m_{00}^2 h^4 s - 1574640 m_{00} g h^3 + 364500 m_{00}^3 h^3 t \\ & + 3458700 m_{00}^4 h^2 s^2 - 2041200 m_{00}^3 g h s + 472500 m_{00}^5 h s t - 122500 m_{00}^6 s^3 + 291600 m_{00}^2 g^2 \\ & - 135000 m_{00}^4 g t + 15625 m_{00}^6 t^2. \end{aligned}$$

This polynomial has 5100 terms in the $m_{i_1 i_2}$.

The moments of order ≤ 3 of probability measures on the triangle $\Delta \subset \mathbb{R}^2$
whose densities are linear functions

The moments of order ≤ 3 of probability measures on the triangle $\Delta \subset \mathbb{R}^2$ whose densities are linear functions form a hypersurface in \mathbb{P}^9 of \mathbb{Z}^3 -degree **(52, 36, 36)**:

The moments of order ≤ 3 of probability measures on the triangle $\triangle \subset \mathbb{R}^2$ whose densities are linear functions form a hypersurface in \mathbb{P}^9 of \mathbb{Z}^3 -degree **(52, 36, 36)**:

$$\begin{aligned}
& 12288754756878336m^{16}s^9 - 125913170530271232h^2m^{14}s^8 - 11555266180939776hm^{15}s^7t - 423695444226048m^{16}s^6t^2 \\
& - 242587475329941504h^4m^{12}s^7 - 67888179490848768h^3m^{13}s^6t - 2253544388296704h^2m^{14}s^5t^2 + 92156256976896hm^{15}s^4t^3 \\
& + 4239929831616m^{16}s^3t^4 - 2425179321925632ghm^{13}s^7 + 767341894828032gm^{14}s^6t - 1302706722212675584h^6m^{10}s^6t^4 \\
& - 108262506929061888h^5m^{11}s^5t + 673312350928896h^4m^{12}s^4t^2 + 535497484271616h^3m^{13}s^3t^3 + 31959518257152h^2m^{14}s^2t^4 \\
& + 440798423040hm^{15}st^5 + 195936798885543936gh^3m^{11}s^6 - 410140620619776gh^2m^{12}s^5t + 412398826108747776gh^6m^8s^3t^2 \\
& - 2360537593675776ghm^{13}s^4t^2 - 89805332054016gm^{14}s^3t^3 - 486870353365172224h^8m^8s^5 + 6819936693387264h^7m^9s^4t \\
& + 29422733985054720h^6m^{10}s^3t^2 + 2782917213290496h^5m^{11}s^2t^3 + 58246341746688h^4m^{12}st^4 - 587731230720h^3m^{13}t^5 \\
& + 3602104581095424g^2m^{12}s^6 - 157746980481662976gh^5m^9s^5 - 79828890012352512gh^4m^{10}s^4t - 10700934975848448gh^3m^{11}s^3t^2 \\
& - 668738492301312gh^2m^{12}s^2t^3 - 10448555212800ghm^{13}st^4 + 275499014400gm^{14}t^5 + 1321196639636946944h^{10}m^6s^4 \\
& + 814698134331457536h^9m^7s^3t + 92179893357379584h^8m^8s^2t^2 + 2541749079638016h^7m^9st^3 - 13792092880896h^6m^{10}t^4 \\
& + 58678654946770944g^2h^2m^{10}s^5 + 16167862146170880g^2hm^{11}s^4t + 705486447968256g^2m^{12}s^3t^2 - 1103687847816200192gh^7m^7s^4 \\
& + 13931406950400gh^3m^{11}t^4 - 44584171418419200gh^5m^9s^2t^2 - 9685512225m^{16}t^6 - 1132386035171328gh^4m^{10}st^3 \\
& + 7839053087502237696h^{12}m^4s^3 + 1352219532013338624h^{11}m^5s^2t + 51427969540816896h^{10}m^6st^2 - 147941222252544h^9m^7t^3 \\
& + 356552602772570112g^2h^4m^8s^4 + 65355404946702336g^2h^3m^9s^3t + 5201278745444352g^2h^2m^{10}s^2t^2 + 99067782758400g^2hm^{11}st^3 \\
& - 3265173504000g^2m^{12}t^4 - 5301992678571900928gh^9m^5s^3 - 984505782412247040gh^8m^6s^2t - 37440870596739072gh^7m^7st^2 \\
& + 260713381625856gh^6m^8t^3 + 7163309458867617792h^{14}m^2s^2 + 495888540219998208h^{13}m^3st - 613682107121664h^{12}m^4t^2 \\
& - 33414364526542848g^3hm^9s^4 - 2441030167166976g^3m^{10}s^3t + 1297818789047435264g^2h^6m^6s^3 + 235088951956733952g^2h^5m^7s^2t \\
& + 8250658482290688g^2h^4m^8st^2 - 132090377011200g^2h^3m^9t^3 - 7123133303988682752gh^{11}m^3s^2 - 506754841838616576gh^{10}m^4st \\
& + 2079004689432576gh^9m^5t^2 + 1846757322198614016h^{16}s - 126388861612851200g^3h^3m^7s^3 - 17847573389770752g^3h^2m^8s^2t \\
& - 469654673817600g^3hm^9st^2 + 20639121408000g^3m^{10}t^3 + 2594242435278176256g^2h^8m^4s^2 + 183620365983940608g^2h^7m^5st \\
& - 1848091141472256g^2h^6m^6t^2 - 2445243491429646336gh^{13}ms + 5610807836540928gh^{12}m^2t + 3143555283419136g^4m^8s^3 \\
& - 408993036765233152g^3h^5m^5s^2 - 26702361435045888g^3h^4m^6st + 626206231756800g^3h^3m^7t^2 + 1246806603479384064g^2h^{10}m^2s \\
& - 9737274975584256g^2h^9m^3t + 22822562857746432g^4h^2m^6s^2 + 1113255523123200g^4hm^7st - 73383542784000g^4m^8t^2 \\
& - 299841218941026304g^3h^7m^3s + 5822326385934336g^3h^6m^4t - 12824703626379264g^2h^{12} + 32389413531025408g^4h^4m^4s \\
& - 1484340697497600g^4h^3m^5t + 15199648742375424g^3h^9m - 1055531162664960g^5hm^5s + 139156940390400g^5m^6t \\
& - 6878544743366656g^4h^6m^2 + 1407374883553280g^5h^3m^3 - 109951162777600g^6m^4.
\end{aligned}$$

Generating Functions

Lemma (Gravin, Pasechnik, Shapiro, Shapiro / Baldoni, Berline, De Loera, Köppe, Vergne)

Let $\Delta_d \subset \mathbb{R}^d$ be a d -dimensional simplex with vertices x_1, x_2, \dots, x_{d+1} .

$$\sum_{\mathcal{I} \in \mathbb{Z}_{\geq 0}^d} \binom{|\mathcal{I}| + d}{\mathcal{I}, d} \cdot m_{\mathcal{I}}(\Delta_d) \cdot t^{\mathcal{I}} =$$

Generating Functions

Lemma (Gravin, Pasechnik, Shapiro, Shapiro / Baldoni, Berline, De Loera, Köppe, Vergne)

Let $\Delta_d \subset \mathbb{R}^d$ be a d -dimensional simplex with vertices x_1, x_2, \dots, x_{d+1} .

$$\sum_{\mathcal{I} \in \mathbb{Z}_{\geq 0}^d} \binom{|\mathcal{I}| + d}{\mathcal{I}, d} \cdot m_{\mathcal{I}}(\Delta_d) \cdot t^{\mathcal{I}} = \prod_{k=1}^{d+1} \frac{1}{1 - \langle x_k, t \rangle}$$

Generating Functions

Lemma (Gravin, Pasechnik, Shapiro, Shapiro / Baldoni, Berline, De Loera, Köppe, Vergne)

Let $\Delta_d \subset \mathbb{R}^d$ be a d -dimensional simplex with vertices x_1, x_2, \dots, x_{d+1} .

$$\sum_{\mathcal{I} \in \mathbb{Z}_{\geq 0}^d} \binom{|\mathcal{I}| + d}{\mathcal{I}, d} \cdot m_{\mathcal{I}}(\Delta_d) \cdot t^{\mathcal{I}} = \prod_{k=1}^{d+1} \frac{1}{1 - \langle x_k, t \rangle}$$

Example ($d = 1$): $\Delta_1 = [a, b] \subset \mathbb{R}^1$

$$\sum_{i=0}^{\infty} (i+1) \cdot m_i \cdot t^i = \frac{1}{(1-at)(1-bt)}$$

Generating Functions

Proposition (K., Shapiro, Sturmfels)

Let $P \subset \mathbb{R}^d$ be a convex polytope with vertices x_1, x_2, \dots, x_n .

$$\sum_{\mathcal{I} \in \mathbb{Z}_{\geq 0}^d} \binom{|\mathcal{I}| + d}{\mathcal{I}, d} \cdot m_{\mathcal{I}}(P) \cdot t^{\mathcal{I}} = \frac{\text{adj}_P(t)}{\prod_{k=1}^n (1 - \langle x_k, t \rangle)}$$

Generating Functions

Proposition (K., Shapiro, Sturmfels)

Let $P \subset \mathbb{R}^d$ be a convex polytope with vertices x_1, x_2, \dots, x_n .

$$\sum_{\mathcal{I} \in \mathbb{Z}_{\geq 0}^d} \binom{|\mathcal{I}| + d}{\mathcal{I}, d} \cdot m_{\mathcal{I}}(P) \cdot t^{\mathcal{I}} = \frac{\text{adj}_P(t)}{\prod_{k=1}^n (1 - \langle x_k, t \rangle)}$$

The **adjoint polynomial** $\text{adj}_P(t)$ of P was introduced by Joe Warren in 1996 to define barycentric coordinates on P .

Generating Functions

Proposition (K., Shapiro, Sturmfels)

Let $P \subset \mathbb{R}^d$ be a convex polytope with vertices x_1, x_2, \dots, x_n .



$$\sum_{\mathcal{I} \in \mathbb{Z}_{\geq 0}^d} \binom{|\mathcal{I}| + d}{\mathcal{I}, d} \cdot m_{\mathcal{I}}(P) \cdot t^{\mathcal{I}} = \frac{\text{adj}_P(t)}{\prod_{k=1}^n (1 - \langle x_k, t \rangle)}$$

The **adjoint polynomial** $\text{adj}_P(t)$ of P was introduced by Joe Warren in 1996 to define barycentric coordinates on P . It was first defined for polygons by Wachspress in 1975:

The Adjoint of a Polygon

Definition (Wachspress)

The **adjoint** A_P of a polygon $P \subset \mathbb{P}^2$ is the minimal degree curve passing through the intersection points of pairs of lines containing non-adjacent edges of P .




$$(\deg A_P = |V(P)| - 3)$$

The Adjoint of a Polygon

Definition (Wachspress)

The **adjoint** A_P of a polygon $P \subset \mathbb{P}^2$ is the minimal degree curve passing through the intersection points of pairs of lines containing non-adjacent edges of P .



$$(\deg A_P = |V(P)| - 3)$$

Proposition (Warren)

Wachspress' adjoint curve A_P of P is defined by Warren's adjoint polynomial $\text{adj}_{P^*}(t)$ of the dual polygon P^* , i.e. $\mathbf{Z}(\text{adj}_{P^*}) = A_P$.

(Recall: $P^* = \{t \in \mathbb{R}^n \mid \forall x \in V(P) : \langle x, t \rangle \leq 1\}$)

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P


The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P




The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P






The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P






The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P






$$\square_1 \cap \square_2 \cap \square_3$$



$$\Delta_1 \cap \Delta_2$$

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



$$\square_1 \cap \square_2 \cap \square_3$$






$$\Delta_1 \cap \Delta_2$$

Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes),

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d **with n facets**
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



$$\square_1 \cap \square_2 \cap \square_3$$






$$\Delta_1 \cap \Delta_2$$

Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes),

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d **with n facets**
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



$$\square_1 \cap \square_2 \cap \square_3$$



$$\Delta_1 \cap \Delta_2$$




Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes), there is a unique hypersurface A_P in \mathbb{P}^d of degree $n - d - 1$ passing through \mathcal{R}_P .

A_P is called the **adjoint** of P .

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d **with n facets**
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



$$\square_1 \cap \square_2 \cap \square_3$$



$$\Delta_1 \cap \Delta_2$$

adjoint plane



Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes), there is a unique hypersurface A_P in \mathbb{P}^d of degree $n - d - 1$ passing through \mathcal{R}_P .


A_P is called the **adjoint** of P .

The Adjoint of a Polytope

- ◆ P : polytope in \mathbb{P}^d **with n facets**
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



adjoint quadric surface



$$\square_1 \cap \square_2 \cap \square_3$$

$$\bullet$$

$$\Delta_1 \cap \Delta_2$$

adjoint plane


Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes), there is a unique hypersurface A_P in \mathbb{P}^d of degree $n - d - 1$ passing through \mathcal{R}_P .


A_P is called the **adjoint** of P .

The Adjoint of a Polytope



- ◆ P : polytope in \mathbb{P}^d **with n facets**
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



adjoint double plane



adjoint quadric surface



adjoint plane


Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes), there is a unique hypersurface A_P in \mathbb{P}^d of degree $n - d - 1$ passing through \mathcal{R}_P .


A_P is called the **adjoint** of P .

The Adjoint of a Polytope



- ◆ P : polytope in \mathbb{P}^d **with n facets**
- ◆ \mathcal{H}_P : hyperplane arrangement spanned by facets of P
- ◆ \mathcal{R}_P : **residual arrangement** of linear spaces that are intersections of hyperplanes in \mathcal{H}_P and do not contain any of face of P



adjoint double plane



adjoint quadric surface




adjoint plane

Theorem (K., Ranestad)

If \mathcal{H}_P is simple (i.e. through any point in \mathbb{P}^d pass $\leq d$ hyperplanes), there is a unique hypersurface A_P in \mathbb{P}^d of degree $n - d - 1$ passing through \mathcal{R}_P .

A_P is called the **adjoint** of P . Moreover, $Z(\text{adj}_{P^*}) = A_P$.



**Thanks for your
attention**

